arXiv:1401.2872v2 [physics.gen-ph] 5 Feb 2014

Hawking radiation - quasi-normal modes correspondence and effective states for nonextremal Reissner-Nordström black holes

[1,2,*]C. Corda, [3,4,**]S. H. Hendi, [5,+]R. Katebi, [6,++]N. O. Schmidt

February 6, 2014

[1]Dipartimento di Fisica e Chimica, Università Santa Rita and Institute for Theoretical Physics and Advanced Mathematics Einstein-Galilei (IFM), 59100 Prato, Italy

[2]International Institute for Applicable Mathematics & Information Sciences (IIAMIS), Hyderabad (India) & Udine (Italy)

[3]Physics Department and Biruni Observatory, College of Sciences, Shiraz University, Shiraz 71454, Iran

[4]Research Institute for Astrophysics and Astronomy of Maragha (RIAAM), P.O. Box 55134-441, Maragha, Iran

[5]Department of Physics, California State University Fullerton, 800 North State College Boulevard, Fullerton, CA 92831, USA

[6]Department of Mathematics, Boise State University, 1910 University Drive, Boise, ID 83725, USA

E-mail addresses: *cordac.galilei@gmail.com, **hendi@shirazu.ac.ir, +rkatebi.gravity@gmail.com, ++nathanschmidt@u.boisestate.edu

Abstract

It is known that the nonstrictly thermal character of the Hawking radiation spectrum harmonizes Hawking radiation with black hole (BH) quasi-normal modes (QNM). This paramount issue has been recently analyzed in the framework of both Schwarzschild BHs (SBH) and Kerr BHs (KBH). In this assignment, we generalize the analysis to the framework

of *nonextremal* Reissner-Nordström BHs (RNBH). Such a generalization is important because in both SBHs and KBHs an absorbed (or emitted) particle has only mass. Instead, in RNBHs the particle has charge as well as mass. In doing so, we expose that for the RNBH, QNMs can be naturally interpreted in terms of quantum levels for both particle emission and absorption. Conjointly, we generalize some concepts concerning the RNBH's "effective states".

1 Introduction

A RNBH of mass M is identical to a SBH of mass M except that a RNBH has the *nonzero* charge quantity Q. In this paper, we are interested in RNBHs with the *nonextremal* constraint $M > Q$ [1]. The quantity Q is the physical mechanism for the RNBH's *dual* horizons from eq. (1) in [1]

$$r_{\pm} = R_{\pm_{RNBH}}(M, Q) = M \pm \sqrt{M^2 - Q^2}, \tag{1}$$

because the RNBH outer (event) horizon radius $R_{+_{RNBH}}(M, Q)$ and the *RNBH inner (Cauchy) horizon radius* $R_{-_{RNBH}}(M, Q)$ are clearly functions of both M *and* Q, not just M as in the well known case of the *SBH horizon radius*

$$r_s = R_{SBH}(M) = 2M. \tag{2}$$

Energy conservation plays a fundamental role in BH radiance [2] because the emission or absorption of a Hawking quanta with mass m and energy-frequency ω causes a BH of mass M to undergo a transition between *discrete* energy spectrum levels [3]-[7], where

$$E = m = \omega = \Delta M \tag{3}$$

for $G = c = k_B = \hbar = \frac{1}{4\pi\epsilon_0} = 1$ (Planck units). Given that emission and absorption are *reverse* processes for the quantized energy spectrum conservation [3]-[7], we consider this pair of transitions as being *equal in magnitude* but *opposite in direction* from the neutral radius perspective of $r_0 = (r_+ + r_-)/2$.

It is known that the countable character of successive emissions of Hawking quanta which is a consequence of the nonstrictly thermal character of the Hawking radiation spectrum (see [3]-[7] and [22]-[26]), generates a natural correspondence between Hawking radiation and BH QNM [3]-[7]. Moreover, it has also been shown that QNMs can be naturally interpreted in terms of quantum levels, where the emission or absorption of a particle is interpreted as a transition between two distinct levels on the *discrete* energy spectrum [3]-[7]. The thermal spectrum correction is an imperative adjustment to the physical interpretation of BH QNMs because these results are important to realize the underlying unitary quantum gravity theory [3]-[7]. Hod's intriguing works [8, 9] suggested that BH QNMs carry principle information regarding a BH's horizon area quantization. Hod's influential conjecture was later refined and clarified by Maggiore [10]. Moreover, it is also believed that QNMs delve into the micro structure of spacetime [11].

To make sense of the state space for the energy spectrum states and the underlying BH perturbation field states, an *effective framework* based on the *nonstrictly thermal* behavior of Hawking's framework began to emerge [3]-[7]. In the midst of this superceding BH effective framework [3]-[7], the BH effective state concept was originally introduced for KBHs in [6] and subsequently applied through Hawking's periodicity arguments [30, 31], to the BH tunneling mechanism's nonstrictly black body spectrum [7]. The effective state is meaningful to BH physics and thermodynamics research because one needs additional features and knowledge to consider in future experiments and observations.

In this paper, our objective is to apply the nonstrictly thermal BH effective framework of [3]-[7] to nonextremal RNBHs. Thus, upon recalling that a RNBH of mass M is identical to a SBH of mass M except that a RNBH has the charge Q, we prepare for our BH QNM investigation by reviewing relevant portions of the SBH effective framework [3]-[7] for quantities related to SBH states and transitions in Section 2. Then in Section 3, we launch our RNBH QNM exploration by introducing a RNBH effective framework for quantities pertaining to RNBH states and transitions. Finally, we conclude with a brief comparison between the fundamental SBH and RNBH results of Section 4 followed by the recapitulation of Section 5.

2 Schwarzschild black hole framework: background and review

2.1 Schwarzschild black hole states and transitions

Here, we recall some quantities that characterize the SBH.

First, consider a SBH of initial mass M, when the SBH emits or absorbs a quantum of energy-frequency ω (for particle mass m and SBH mass change ΔM, such that $m = \omega = \Delta M$) to achieve a final mass of $M - \omega$ or $M + \omega$, respectively, for the SBH mass-energy transition between states in state space. Thus, we follow [3]-[5], where the *SBH initial and final horizon area* are

$$\begin{aligned} A_{SBH}(M) &= 16\pi M^2 &&= 4\pi R_{SBH}^2(M) \\ A_{SBH}(M \pm \omega) &= 16\pi (M \pm \omega)^2 &&= 4\pi R_{SBH}^2(M \pm \omega), \end{aligned} \tag{4}$$

respectively, for the *SBH area quanta number*

$$N_{SBH}(M, \omega) = \frac{A_{SBH}(M)}{|\Delta A_{SBH}(M, \omega)|}, \tag{5}$$

such that the *SBH horizon area change* for the corresponding mass change ΔM is

$$\begin{aligned} \Delta A_{SBH}(M, \omega) &= A_{SBH}(M \pm \omega) - A_{SBH}(M) \\ &= 32\pi M\omega + O(\omega^2) \sim 32\pi M \Delta M \\ &= 32\pi M \Delta E, \end{aligned} \tag{6}$$

because the transition's minus $(-)$ and plus $(+)$ signs depend on emission and absorption, respectively. Next, in [3]-[5], the *Bekenstein-Hawking SBH initial and final entropy* are

$$\begin{aligned} S_{SBH}(M) &= \tfrac{A_{SBH}(M)}{4} \\ S_{SBH}(M \pm \omega) &= \tfrac{A_{SBH}(M \pm \omega)}{4}, \end{aligned} \tag{7}$$

respectively, where the corresponding *SBH entropy change* is

$$\Delta S_{SBH}(M, \omega) = \frac{\Delta A_{SBH}(M, \omega)}{4}. \tag{8}$$

Subsequently, the *SBH initial and final total entropy* are [3]-[5]

$$\begin{aligned} S_{SBH-total}(M) &= S_{SBH}(M) - \ln S_{SBH}(M) + \tfrac{3}{2A_{SBH}(M)} \\ S_{SBH-total}(M \pm \omega) &= S_{SBH}(M \pm \omega) - \ln S_{SBH}(M \pm \omega) + \tfrac{3}{2A_{SBH}(M \pm \omega)}, \end{aligned} \tag{9}$$

respectively. Additionally, the *SBH initial and final Hawking temperature* are [3]-[5]

$$\begin{aligned} T_{H_{SBH}}(M) &= \tfrac{1}{8\pi M} \\ T_{H_{SBH}}(M \pm \omega) &= \tfrac{1}{8\pi (M \pm \omega)}, \end{aligned} \tag{10}$$

respectively. Therefore, the quantum transition's *SBH emission tunneling rate* is [3]-[5]

$$\begin{aligned} \Gamma_{SBH}(M, \omega) &\sim \exp\left[-8\pi M\omega \left(1 - \tfrac{\omega}{2M}\right)\right] \\ &\sim \exp\left[-\tfrac{\omega}{T_{H_{SBH}}(M)} \left(1 - \tfrac{\omega}{R_{SBH}(M)}\right)\right] \\ &\sim \exp\left[+\Delta S_{SBH}(M, \omega)\right]. \end{aligned} \tag{11}$$

2.2 Schwarzschild black hole effective states and transitions

Here, we recall some effective quantities that characterize the SBH.

Given that M is the mass state *before* and $M \pm \omega$ is the mass state *after* the quantum transition, the *SBH effective mass* and *SBH effective horizon* are respectively identified in [3]-[5] as

$$\begin{aligned} M_E(M, \omega) &= \tfrac{M + (M \pm \omega)}{2} = M \pm \tfrac{\omega}{2} \\ R_{E_{SBH}}(M, \omega) &= 2M_E(M, \omega) \end{aligned}, \tag{12}$$

which are *average quantities* between the two states *before* and *after* the process [3]-[5]. Consequently, using eqs. (4) and (12) we define the *SBH effective horizon area* as

$$\begin{aligned} A_{E_{SBH}}(M, \omega) \equiv \tfrac{A_{SBH}(M) + A_{SBH}(M \pm \omega)}{2} = \\ \equiv 16\pi M_E^2(M, \omega) = 4\pi R_{E_{SBH}}^2(M, \omega), \end{aligned} \tag{13}$$

which is the average of the SBH's initial and final horizon areas. Subsequently, utilizing eq. (7), the Bekenstein-Hawking *SBH effective entropy* is defined as

$$S_{E_{SBH}}(M,\omega) \equiv \frac{S_{SBH}(M) + S_{SBH}(M \pm \omega)}{2}, \tag{14}$$

and consequently employs eqs. (13) and (14) to define the *SBH effective total entropy* as

$$\begin{aligned} & S_{E_{SBH-total}}(M,\omega) \equiv \\ & \equiv S_{E_{SBH}}(M,\omega) - \ln S_{E_{SBH}}(M,\omega) + \tfrac{3}{2A_{E_{SBH}}(M,\omega)}. \end{aligned} \tag{15}$$

Thus, employing eqs. (3) and (10), the *SBH effective temperature* is [3]-[5]

$$\begin{aligned} T_{E_{SBH}}(M,\omega) &= \left(\tfrac{T^{-1}_{H_{SBH}}(M)+T^{-1}_{H_{SBH}}(M\pm\omega)}{2}\right)^{-1} \\ &= \left(8\pi\left[\tfrac{M+M\pm\omega}{2}\right]\right)^{-1} \\ &= \tfrac{1}{4\pi(2M\pm\omega)} = \tfrac{1}{8\pi M_E(M,\omega)}, \end{aligned} \tag{16}$$

which is the inverse of the average value of the inverses of the initial and final Hawking temperatures. Consequently, eq. (16) lets one rewrite eq. (11) to define the *SBH effective emission tunneling rate* (in the Boltzmann-like form) as [3]-[5]

$$\Gamma_{E_{SBH}}(M,\omega) \sim \exp\left[-\frac{\omega}{T_{E_{SBH}}(M,\omega)}\right] = \exp\left[+\Delta S_{E_{SBH}}(M,\omega)\right], \tag{17}$$

such that eq. (14) defines the *SBH effective entropy change* as

$$\Delta S_{E_{SBH}}(M,\omega) = S_{SBH}(M \pm \omega) - S_{SBH}(M) = \frac{\Delta A_{E_{SBH}}(M,\omega)}{4} \tag{18}$$

because the *SBH effective horizon area change* is

$$\Delta A_{E_{SBH}}(M,\omega) = 16\pi M_E(M,\omega)\omega \tag{19}$$

and the *SBH effective area quanta number* is

$$N_{E_{SBH}}(M,\omega) = \frac{A_{E_{SBH}}(M,\omega)}{\Delta A_{E_{SBH}}(M,\omega)}. \tag{20}$$

2.3 Effective application of quasi-normal modes to the Schwarzschild black hole

Here, we recall how the SBH perturbation field QNM states can be applied to the SBH effective framework.

The quasi-normal frequencies (QNF) are typically labeled as ω_{nl}, where l is the angular momentum quantum number [3]-[5, 10, 12]. Thus, for each l,

such that $l \geq 2$ for gravitational perturbations, there is a countable sequence of QNMs labeled by the overtone number n, which is a natural number [3]-[5, 10].

Now $|\omega_n|$ is the damped harmonic oscillator's proper frequency that is defined as [3]-[5, 10]

$$|\omega_n| = (\omega_0)_n = \sqrt{\omega_{n_{\mathbb{R}}}^2 + \omega_{n_{\mathbb{I}}}^2}. \tag{21}$$

Maggiore [10] articulated that the establishment $|\omega_n| = \omega_{n_{\mathbb{R}}}$ is only correct for the very long-lived and lowly excited QNMs approximation $|\omega_n| \gg \omega_{n_{\mathbb{I}}}$, whereas for a lot of BH QNMs, such as those that are highly excited, the opposite limit is correct [3]-[5, 10]. Therefore, the ω parameter in eqs. $(12) - (20)$ is substituted for the $|\omega_n|$ parameter [3]-[5] because we wish to employ BH QNFs. When n is large the SBH QNFs become independent of l and thereby exhibit the nonstrictly thermal structure [3]-[5]

$$\begin{aligned}
\omega_n &= \ln 3 \times T_{E_{SBH}}(M, |\omega_n|) + 2\pi i(n + \tfrac{1}{2}) \times T_{E_{SBH}}(M, |\omega_n|) + \mathcal{O}(n^{-\frac{1}{2}}) \\
&= \tfrac{\ln 3}{4\pi[2M - |\omega_n|]} + \tfrac{2\pi i}{4\pi[2M - |\omega_n|]}(n + \tfrac{1}{2}) + \mathcal{O}(n^{-\frac{1}{2}}) \\
&= \tfrac{\ln 3}{8\pi M_E(M, |\omega_n|)} + \tfrac{2\pi(n+\frac{1}{2})}{8\pi M_E(M, |\omega_n|)} i + \mathcal{O}(n^{-\frac{1}{2}}),
\end{aligned} \tag{22}$$

where

$$\begin{aligned}
m_n \equiv \omega_{n_{\mathbb{R}}} &= \tfrac{\ln 3}{8\pi M_E(M, |\omega_n|)} \\
p_n \equiv \omega_{n_{\mathbb{I}}} &= \tfrac{2\pi}{8\pi M_E(M, |\omega_n|)}(n + \tfrac{1}{2}).
\end{aligned} \tag{23}$$

Thus, when referring to highly excited QNMs one gets $|\omega_n| \approx p_n$ [3]-[5], where the quantized levels differ from [10] because they are not equally spaced in exact form. Therefore, according to [3]-[5], we have

$$\begin{aligned}
|\omega_n| &= \tfrac{\sqrt{(\ln 3)^2 + 4\pi^2(n+\frac{1}{2})^2}}{8\pi M_E(M, |\omega_n|)} \\
&= T_{E_{SBH}}(M, |\omega_n|)\sqrt{(\ln 3)^2 + 4\pi^2(n + \tfrac{1}{2})^2},
\end{aligned} \tag{24}$$

which is solved to yield

$$|\omega_n| = M - \sqrt{M^2 - \frac{\sqrt{(\ln 3)^2 + 4\pi^2(n + \frac{1}{2})^2}}{4\pi}} \tag{25}$$

when we obey $|\omega_n| < M$ because a BH cannot emit more energy than its total mass.

3 Reissner-Nordström black hole framework: an introduction

We note that for this framework, we consider the *RNBH event horizon* features, which are derived from the $R_{+_{RNBH}}(M, Q)$ in eq. (1).

3.1 Reissner-Nordström black hole states and transitions

Here, we recall some quantities that characterize the RNBH.

First, consider a RNBH of initial mass M and initial charge Q. Using eq. (1), we define the *RNBH initial event horizon area* as

$$A_{+_{RNBH}}(M,Q) = 4\pi(M+\sqrt{M^2-Q^2})^2 = 4\pi R^2_{+_{RNBH}}(M,Q), \tag{26}$$

the *Bekenstein-Hawking RNBH initial entropy* as

$$S_{+_{RNBH}}(M,Q) = \frac{A_{+_{RNBH}}(M,Q)}{4}, \tag{27}$$

and the *RNBH initial electrostatic potential* as

$$\Phi_{+}(M,Q) = \frac{Q}{4\pi R_{+_{RNBH}}(M,Q)} = \frac{Q}{4\pi(M+\sqrt{M^2-Q^2})}. \tag{28}$$

Consequently, eq. (17) of [2] identifies the *RNBH initial Hawking temperature* as

$$\begin{aligned} T_{+H_{RNBH}}(M,Q) &= \frac{\sqrt{M^2-Q^2}}{2\pi(M+\sqrt{M^2-Q^2})^2} = \\ &= \frac{R_{+_{RNBH}}(M,Q)-R_{-_{RNBH}}(M,Q)}{A_{+RNBH}(M,Q)}. \end{aligned} \tag{29}$$

Second, consider when the RNBH emits or absorbs a quantum of energy-frequency ω with charge q to achieve a final mass of $M-\omega$ or $M+\omega$ and a final charge of $Q-q$ or $Q+q$, respectively, for the RNBH mass-energy transition between states in state space. For this, all we need to do is replace the RNBH's mass and charge parameters in eqs. (26) and (29). Thus, eq. (26) establishes the *RNBH final event horizon area* as

$$\begin{aligned} A_{+_{RNBH}}(M\pm\omega,Q\pm q) &= 4\pi R^2_{+_{RNBH}}(M\pm\omega,Q\pm q) \\ &= 4\pi\left((M\pm\omega)+\sqrt{(M\pm\omega)^2-(Q\pm q)^2}\right)^2, \end{aligned} \tag{30}$$

eq. (27) presents the *Bekenstein-Hawking RNBH final entropy* as

$$S_{+_{RNBH}}(M\pm\omega,Q\pm q) = \frac{A_{+_{RNBH}}(M\pm\omega,Q\pm q)}{4}, \tag{31}$$

and eq. (28) defines the *RNBH final electrostatic potential* as

$$\begin{aligned} \Phi_{+}(M\pm\omega,Q\pm q) &= \frac{Q}{4\pi R_{+_{RNBH}}(M\pm\omega,Q\pm q)} \\ &= \frac{Q}{4\pi\left((M\pm\omega)+\sqrt{(M\pm\omega)^2-(Q\pm q)^2}\right)} \end{aligned} \tag{32}$$

for usage in eq. (29) of [13], where it is proposed that the *RNBH adiabatic invariant* is

$$\begin{aligned} I_{+_{RNBH}}(M,\omega,Q,q) = \int \frac{\omega - \Phi_+(M\pm\omega,Q)q}{\omega} = \\ = \int \frac{\Delta M - \Phi_+(M\pm\Delta M,Q)\Delta Q}{\Delta M} \end{aligned} \tag{33}$$

because $\Delta Q = q$. Hence, eq. (29) identifies the *RNBH final Hawking temperature* as

$$\begin{aligned} T_{+H_{RNBH}}(M \pm \omega, Q \pm q) &= \frac{R_{+_{RNBH}}(M\pm\omega,Q\pm q) - R_{-_{RNBH}}(M\pm\omega,Q\pm q)}{A_{+RNBH}(M\pm\omega,Q\pm q)} \\ &= \frac{\sqrt{(M\pm\omega)^2-(Q\pm q)^2}}{2\pi\left((M\pm\omega)+\sqrt{(M\pm\omega)^2-(Q\pm q)^2}\right)^2}. \end{aligned} \tag{34}$$

Next, upon generalizing eq. (16) in [2] and the work [27], we define the *RNBH tunneling rate* as

$$\begin{aligned} \Gamma_{+_{RNBH}}(M,\omega,Q,q) \sim \\ \exp\left[-4\pi\left(2\omega(M\pm\tfrac{\omega}{2}) - (M\pm\omega)\sqrt{(M\pm\omega)^2-(Q\pm q)^2} + M\sqrt{M^2-Q^2}\right)\right] \sim \\ \exp\left[\Delta S_{+_{RNBH}}(M,\omega,Q,q)\right], \end{aligned} \tag{35}$$

where we utilize eq. (30) to define the *Bekenstein-Hawking RNBH entropy change* as

$$\Delta S_{+_{RNBH}}(M,\omega,Q,q) = \frac{\Delta A_{+RNBH}(M,\omega,Q,q)}{4}, \tag{36}$$

such that the *RNBH event horizon area change* is

$$\Delta A_{+_{RNBH}}(M,\omega,Q,q) = A_{+_{RNBH}}(M\pm\omega,Q\pm q) - A_{+_{RNBH}}(M,Q) \tag{37}$$

so we can define the *RNBH event horizon area quanta number* as

$$N_{+_{RNBH}}(M,\omega,Q,q) = \frac{A_{+_{RNBH}}(M,Q)}{|\Delta A_{+_{RNBH}}(M,\omega,Q,q)|}. \tag{38}$$

3.2 Reissner-Nordström black hole effective states and transitions

Here, we define some effective quantities that characterize the RNBH.

The *RNBH effective mass* is equivalent to the SBH effective mass component of eq. (12), which is

$$M_E(M,\omega) \equiv \frac{M + (M\pm\omega)}{2}. \tag{39}$$

Next, we define the *RNBH effective charge* as

$$Q_E(Q,q) \equiv \frac{Q + (Q\pm q)}{2}, \tag{40}$$

which is the average of the RNBH's initial charge Q and final charge $Q \pm q$. From this, eqs. (1), (39) and (40) are used to define the corresponding *RNBH effective event horizon* and *RNBH effective cauchy horizon* as

$$r_{\pm E} \equiv R_{\pm E_{RNBH}}(M,\omega,Q,q) \equiv M_E(M,\omega) \pm \sqrt{M_E^2(M,\omega) - Q_E^2(Q,q)}, \quad (41)$$

with respect to the energy conservation and pair production neutrality of eq. (39). Next, we employ eqs. (26), (39) and (41) to define the *RNBH effective event horizon area* as

$$\begin{aligned} A_{+E_{RNBH}}(M,\omega,Q,q) &\equiv 4\pi R^2_{+E_{RNBH}}(M,\omega,Q,q) \\ &\equiv 4\pi \left(M_E(M,\omega) + \sqrt{M_E^2(M,\omega) - Q_E^2(Q,q)} \right)^2, \end{aligned} \quad (42)$$

which is then used to define the *RNBH effective entropy* as

$$S_{+E_{RNBH}}(M,\omega,Q,q) \equiv \frac{A_{+E_{RNBH}}(M,\omega,Q,q)}{4}. \quad (43)$$

Afterwards, we use eqs. (28) and (42) to define the *RNBH effective electrostatic potential* as

$$\begin{aligned} \Phi_{+E}(M,\omega,Q,q) &\equiv \frac{Q_E(Q,q)}{4\pi R_{+E_{RNBH}}(M,\omega,Q,q)} \\ &\equiv \frac{Q_E(Q,q)}{4\pi \left(M_E(M,\omega) + \sqrt{M_E^2(M,\omega) - Q_E^2(Q,q)} \right)} \end{aligned} \quad (44)$$

so we can utilize the $T_{E_{SBH}}(M,\omega)$ in eq. (16) along with eqs. (39), (40), and (44) to define the *RNBH effective adiabatic invariant* as

$$I_{+E_{RNBH}}(M,\omega,Q,q) \equiv \int \frac{dM_E(M,\omega) - \Phi_{+E}(M,\omega,Q,q) dQ_E(Q,q)}{T_{E_{SBH}}(M,\omega)}. \quad (45)$$

At this point, eqs. (16) and (35) let us introduce and define the *RNBH effective temperature* as

$$\begin{aligned} T_{+E_{RNBH}}(M,\omega,Q,q) &\equiv \frac{\sqrt{\left(M \pm \frac{\omega}{2}\right)^2 - \left(Q \pm \frac{q}{2}\right)^2}}{2\pi \left[\left(M \pm \frac{\omega}{2}\right) + \sqrt{\left(M \pm \frac{\omega}{2}\right)^2 - \left(Q \pm \frac{q}{2}\right)^2} \right]^2} \\ &\equiv \frac{\sqrt{M_E^2(M,\omega) - Q_E^2(Q,q)}}{2\pi \left(M_E(M,\omega) + \sqrt{M_E^2(M,\omega) - Q_E^2(Q,q)} \right)^2} \\ &\equiv \frac{R_{+E_{RNBH}}(M,\omega,Q,q) - R_{-E_{RNBH}}(M,\omega,Q,q)}{A_{+E_{RNBH}}(M,\omega,Q,q)}, \end{aligned} \quad (46)$$

which authorizes us to exercise eqs. (36) and (46) to rewrite eq. (35) to define the RNBH effective tunneling rate as

$$\begin{aligned} \Gamma_{+E_{RNBH}}(M,\omega,Q,q) &\sim \exp\left[\frac{\pm\omega}{T_{+E_{RNBH}}(M,\omega,Q,q)} \right] \\ &\sim \exp\left[\Delta S_{+_{RNBH}}(M,\omega,Q,q) \right], \end{aligned} \quad (47)$$

such that the *RNBH effective entropy change* is defined as

$$\Delta S_{+E_{RNBH}}(M,\omega,Q,q) \equiv \frac{\Delta A_{+E_{RNBH}}(M,\omega,Q,q)}{4} \tag{48}$$

for the *RNBH effective event horizon area change*

$$\Delta A_{+E_{RNBH}}(M,\omega,Q,q) \equiv \frac{2\omega q + Q^3\pi}{(M^2-Q^2)^{3/2}} \tag{49}$$

and the *RNBH effective event horizon area quanta number*

$$N_{+E_{RNBH}}(M,\omega,Q,q) \equiv \frac{A_{+E_{RNBH}}(M,\omega,Q,q)}{|\Delta A_{+E_{RNBH}}(M,\omega,Q,q)|}. \tag{50}$$

4 Effective application of quasi-normal modes to the Reissner-Nordström black hole

Here, we explain how the RNBH perturbation field QNM states can be applied to the RNBH effective framework.

Similarly to SBH QNFs, the RNBH QNFs become independent of l for large n [14]. Thus, for large n, we have two families of the QNM:

$$\omega_n = \ln 3 \times T_{+H_{SBH}}(M,Q) - 2\pi(n+\frac{1}{2})i \times T_{+H_{SBH}}(M,Q) + \frac{qQ}{R_{+_{SBH}}(M,Q)} \tag{51}$$

and

$$\begin{aligned} \omega_n &= \ln 2 \times T_{+H_{RNBH}}(M,Q) - 2\pi(n+\tfrac{1}{2})i \times T_{+H_{RNBH}}(M,Q) + \tfrac{qQ}{R_{+_{RNBH}}(M,Q)} \\ &= \frac{\ln 2\sqrt{M^2-Q^2}}{2\pi\left(M+\sqrt{M^2-Q^2}\right)^2} - \frac{(n+\frac{1}{2})\sqrt{M^2-Q^2}}{\left(M+\sqrt{M^2-Q^2}\right)^2}i + \frac{qQ}{R_{+_{RNBH}}(M,Q)}. \end{aligned} \tag{52}$$

Now the approximation of eqs. (51) and (52) are only relevant under the assumption that the BH radiation spectrum is strictly thermal [3]-[5] because they both use the Hawking temperature $T_{+H_{RNBH}}$ in eq. (29). Hence, to operate in compliance of [3]-[5] and thereby account for the thermal spectrum deviation of eq. (35), we opt to select the eq. (52) case and upgrade it by effectively replacing its $T_{H_{RNBH}}$ in eq. (29) with the $T_{+E_{RNBH}}$ in eq. (46). Therefore, the corrected expression for the RNBH QNFs of eq. (52) that encodes the nonstrictly thermal behavior of the radiation spectrum is defined as

$$\begin{aligned} \omega_n &\equiv \ln 2 \times T_{+E_{RNBH}}(M,|\omega_n|,Q,q) - 2\pi(n+\tfrac{1}{2})i \times T_{+E_{RNBH}}(M,|\omega_n|,Q,q) \\ &\quad + \tfrac{qQ_E(Q,q)}{R_{+E_{RNBH}}(M,|\omega_n|,Q,q)} \\ &\equiv \frac{\ln 2\sqrt{M_E^2(M,|\omega_n|)-Q_E^2(Q,q)}}{2\pi\left(M_E(M,|\omega_n|)+\sqrt{M_E^2(M,|\omega_n|)-Q_E^2(Q,q)}\right)^2} - \frac{(n+\frac{1}{2})\sqrt{M_E^2(M,|\omega_n|)-Q_E^2(Q,q)}}{\left(M_E(M,|\omega_n|)+\sqrt{M_E^2(M,|\omega_n|)-Q_E^2(Q,q)}\right)^2}i \\ &\quad + \frac{qQ_E(Q,q)}{R_{+E_{RNBH}}(M,|\omega_n|,Q,q)}. \end{aligned} \tag{53}$$

From eqs. (39), (41), and (46) we define the effective quantities associated to the QNMs as

$$M_E(M,|\omega_n|) \equiv \frac{M+(M-|\omega_n|)}{2}, \tag{54}$$

$$\begin{aligned} r_{\pm E} &\equiv R_{\pm E_{RNBH}}(M,|\omega_n|,Q,q) \\ &\equiv M_E(M,|\omega_n|) \pm \sqrt{M_E^2(M,|\omega_n|)-Q_E^2(Q,q)}, \end{aligned} \tag{55}$$

and

$$\begin{aligned} T_{+E_{RNBH}}(M,|\omega_n|,Q,q) &\equiv \frac{\sqrt{\left(M-\frac{|\omega_n|}{2}\right)^2-\left(Q-\frac{q}{2}\right)^2}}{2\pi\left[\left(M-\frac{|\omega_n|}{2}\right)+\sqrt{\left(M-\frac{|\omega_n|}{2}\right)^2-\left(Q-\frac{q}{2}\right)^2}\right]^2} \\ &= \frac{\sqrt{M_E^2(M,|\omega_n|)-Q_E^2(Q,q)}}{2\pi\left(M_E(M,|\omega_n|)+\sqrt{M_E^2(M,|\omega_n|)-Q_E^2(Q,q)}\right)^2} \\ &= \frac{R_{+E_{RNBH}}(M,|\omega_n|,Q,q)-R_{-E_{RNBH}}(M,|\omega_n|,Q,q)}{A_{+E_{RNBH}}(M,|\omega_n|,Q,q)}, \end{aligned} \tag{56}$$

respectively, for the quantum overtone number n in eq. (53). Hence, eq. (53) lets us rewrite the SBH case of eq. (23) to present the RNBH case

$$\begin{aligned} m_n &\equiv \ln 2 \times T_{+E_{RNBH}}(M,|\omega_n|,Q,q) + \frac{eQ_E(Q,q)}{R_{+E_{RNBH}}(M,|\omega_n|,Q,q)} \\ &= \frac{\ln 2\sqrt{M_E^2(M,|\omega_n|)-Q_E^2(Q,q)}}{2\pi\left(M_E(M,|\omega_n|)+\sqrt{M_E^2(M,|\omega_n|)-Q_E^2(Q,q)}\right)^2} + \frac{qQ_E(Q,q)}{R_{+E_{RNBH}}(M,|\omega_n|,Q,q)} \end{aligned}$$

$$p_n \equiv -2\pi(n+\tfrac{1}{2}) \times T_{+E_{RNBH}}(M,|\omega_n|,Q,q) = -\frac{(n+\frac{1}{2})\sqrt{M_E^2(M,|\omega_n|)-Q_E^2(Q,q)}}{\left(M_E(M,|\omega_n|)+\sqrt{M_E^2(M,|\omega_n|)-Q_E^2(Q,q)}\right)^2}. \tag{57}$$

Thus, we recall that if $|\omega_n| \approx p_n$, then we are referring to highly excited QNMs [3]-[5] . Therefore, the SBH case of eq. (24) becomes the RNBH case

$$\begin{aligned} |\omega_n| &\equiv \frac{\sqrt{M_E^2(M,|\omega_n|)-Q_E^2(Q,q)}\sqrt{(\ln 2)^2-4\pi^2(n+\frac{1}{2})^2}}{2\pi\left(M_E(M,|\omega_n|)+\sqrt{M_E^2(M,|\omega_n|)-Q_E^2(Q,q)}\right)^2} + \frac{qQ_E(Q,q)}{R_{+E_{RNBH}}(M,|\omega_n|,Q,q)} \\ &= T_{+E_{RNBH}}(M,|\omega_n|,Q,q)\sqrt{(\ln 2)^2-4\pi^2(n+\tfrac{1}{2})^2} + \frac{qQ_E(Q,q)}{R_{+E_{RNBH}}(M,|\omega_n|,Q,q)}. \end{aligned} \tag{58}$$

Hence, upon considering eqs. (40) and (54), one can rewrite eq. (58) as

$$\begin{aligned}|\omega_n| &\equiv \frac{\sqrt{\left(M-\frac{|\omega_n|}{2}\right)^2-\left(Q-\frac{q}{2}\right)^2}\sqrt{(\ln 2)^2-4\pi^2(n+\frac{1}{2})^2}}{2\pi\left[\left(M-\frac{|\omega_n|}{2}\right)+\sqrt{\left(M-\frac{|\omega_n|}{2}\right)^2-\left(Q-\frac{q}{2}\right)^2}\right]^2} \\ &+\frac{q\left(Q-\frac{q}{2}\right)}{\left(M-\frac{|\omega_n|}{2}\right)+\sqrt{\left(M-\frac{|\omega_n|}{2}\right)^2-\left(Q-\frac{q}{2}\right)^2}},\end{aligned} \tag{59}$$

where the solution of eq. (59) in terms of $|\omega_n|$ will be the answer of $|\omega_n|$. Therefore, given a quantum transition between the levels n and $n-1$, we define $|\triangle\omega_{n,n-1}| \equiv |\omega_n - \omega_{n-1}|$ where eqs. (41)-(45) are rewritten as

$$\begin{aligned}r_{\pm E} \equiv R_{\pm E_{RNBH}}(M,|\triangle\omega_{n,n-1}|,Q,q) \equiv \\ \equiv M_E(M,|\triangle\omega_{n,n-1}|) \pm \sqrt{M_E^2(M,|\triangle\omega_{n,n-1}|)-Q_E^2(Q,q)},\end{aligned} \tag{60}$$

$$\begin{aligned}A_{+E_{RNBH}}(M,|\triangle\omega_{n,n-1}|,Q,q) \equiv 4\pi R^2_{+E_{RNBH}}(M,|\triangle\omega_{n,n-1}|,Q,q) \\ \equiv 4\pi\left(M_E(M,|\triangle\omega_{n,n-1}|)+\sqrt{M_E^2(M,|\triangle\omega_{n,n-1}|)-Q_E^2(Q,q)}\right)^2,\end{aligned} \tag{61}$$

$$S_{+E_{RNBH}}(M,|\triangle\omega_{n,n-1}|,Q,q) \equiv \frac{A_{+E_{RNBH}}(M,|\triangle\omega_{n,n-1}|,Q,q)}{4}, \tag{62}$$

$$\begin{aligned}\Phi_{+E}(M,|\triangle\omega_{n,n-1}|,Q,q) &\equiv \frac{Q_E(Q,q)}{4\pi R_{+E_{RNBH}}(M,|\triangle\omega_{n,n-1}|,Q,q)} \\ &\equiv \frac{Q_E(Q,q)}{4\pi\left(M_E(M,|\triangle\omega_{n,n-1}|)+\sqrt{M_E^2(M,|\triangle\omega_{n,n-1}|)-Q_E^2(Q,q)}\right)},\end{aligned} \tag{63}$$

$$\begin{aligned}I_{+E_{RNBH}}(M,|\triangle\omega_{n,n-1}|,Q,q) = \\ = \int \frac{dM_E(M,|\triangle\omega_{n,n-1}|)-\Phi_{+E}(M,|\triangle\omega_{n,n-1}|,Q,q)dQ_E(Q,q)}{T_{E_{SBH}}(M,|\triangle\omega_{n,n-1}|)},\end{aligned} \tag{64}$$

and eqs. (47)-(50) become

$$\begin{aligned}\Gamma_{+E_{RNBH}}(M,|\triangle\omega_{n,n-1}|,Q,q) &\sim \exp\left[\frac{\pm|\triangle\omega_{n,n-1}|}{T_{+E_{RNBH}}(M,|\triangle\omega_{n,n-1}|,Q,q)}\right] \\ &\sim \exp\left[\Delta S_{+_{RNBH}}(M,|\triangle\omega_{n,n-1}|,Q,q)\right],\end{aligned} \tag{65}$$

$$\Delta S_{+E_{RNBH}}(M, |\triangle\omega_{n,n-1}|, Q, q) \equiv \frac{\Delta A_{+E_{RNBH}}(M, |\triangle\omega_{n,n-1}|, Q, q)}{4}, \quad (66)$$

$$\Delta A_{+E_{RNBH}}(M, |\triangle\omega_{n,n-1}|, Q, q) \equiv \frac{2|\triangle\omega_{n,n-1}|q + \pi Q^3}{(M^2 - Q^2)^{3/2}}, \quad (67)$$

$$N_{+E_{RNBH}}(M, |\triangle\omega_{n,n-1}|, Q, q) \equiv \frac{A_{+E_{RNBH}}(M, |\triangle\omega_{n,n-1}|, Q, q)}{|\Delta A_{+E_{RNBH}}(M, |\triangle\omega_{n,n-1}|, Q, q)|}, \quad (68)$$

respectively.

5 A brief comparison

Here, we will show that the SBH results of Section 2 are in fundamental agreement with the RNBH results of Section 3 for small Q, where we recall that the RNBH of mass M is identical to a SBH of mass M except that a RNBH has the nonzero charge quantity Q.

First, for small Q, the SBH's $T_{E_{SBH}}(M, \omega)$ of eq. (16) is related to the RNBH's $T_{+E_{RNBH}}(M, \omega, Q, q)$ of eq. (46) as

$$T_{+E_{RNBH}}(M, \omega, Q, q) \equiv T_{E_{SBH}}(M, \omega) - \frac{3q^2Q^2}{8(2m \pm \omega)^5\pi} + \mathcal{O}(Q^4, q^4). \quad (69)$$

Second, for small Q, the SBH's $A_{E_{SBH}}(M, \omega)$ of eq. (13) complies with the RNBH's $A_{+E_{RNBH}}(M, \omega, Q, q)$ of eq. (42) as

$$A_{+E_{RNBH}}(M, \omega, Q, q) \equiv A_{E_{SBH}}(M, \omega) - 8\pi Q^2 + \mathcal{O}(Q^4). \quad (70)$$

Third, for small Q, the SBH's $S_{E_{SBH}}(M, \omega)$ of eq. (14) corresponds with the RNBH's $S_{+E_{RNBH}}(M, \omega, Q, q)$ of eq. (43) as

$$S_{+E_{RNBH}}(M, \omega, Q, q) \equiv S_{E_{SBH}}(M, \omega) - 2\pi Q^2 + \mathcal{O}(Q^4). \quad (71)$$

Fourth, for small Q, the SBH's QNF $|\omega_n|$ of eq. (24) is consistent with the RNBH's QNF $|\omega_n|$ of eqs. (58) and (59) as

$$\begin{aligned} |\omega_n| &\equiv \tfrac{\sqrt{\ln 2^2 - 4\pi^2(n+\frac{1}{2})^2}}{4(2M-|\omega_n|)\pi} + \tfrac{qQ}{2M-|\omega_n|} = \\ &\tfrac{3(16\pi M^2 - 16\pi|\omega_n|M + 4\pi|\omega_n|^2 + \sqrt{-(\ln 2+\pi+2\pi n)(-\ln 2+\pi+2\pi n)})}{8(2M-|\omega_n|)^5\pi} Q^2q^2 \quad = \\ &\tfrac{q^2}{4M-|\omega_n|} + \mathcal{O}(Q^4, q^4), \end{aligned} \quad (72)$$

which can be applied to eqs. (69)-(70) by replacing the ω parameter with the pertinent $|\omega_n|$. Hence, eqs. (69)-(72) indicate that in general, the SBH results of Section 2 are fundamentally consistent with the RNBH results of Section 3

for small Q. Moreover, in eq. (72) for large n, the result is consistent with the SBH because $\ln 2$ is negligible, but for small n there is an argument between scientists regarding $\ln 2$ and $\ln 3$ because these refer to the two distinct QNM families of eqs. (51) and (52).

Here, we provide the physical answer of eq. (72) for the case of emission by using the fact that Q is small, so the term which includes Q^2 is also very small and therefore negligible:

$$(\omega_0)_n \equiv |\omega_n| \approx M - \sqrt{M^2 + \frac{q^2}{2} - Qq - \frac{1}{4\pi}\sqrt{\ln 2^2 - 4\pi^2(n+\frac{1}{2})^2}}. \qquad (73)$$

Thus, by setting $(\omega_0)_n \equiv |\omega_n|$ we obtain

$$\begin{aligned} \triangle M_n \quad \equiv -\triangle\omega_{n,n-1} = \quad & (\omega_0)_{n-1} - (\omega_0)_n \\ \equiv \quad & \sqrt{M^2 + \tfrac{q^2}{2} - Qq - \tfrac{1}{4\pi}\sqrt{(\ln 2)^2 + 4\pi^2(n+\tfrac{1}{2})^2}} \\ & -\sqrt{M^2 + \tfrac{q^2}{2} - Qq - \tfrac{1}{4\pi}\sqrt{(\ln 2)^2 + 4\pi^2(n-\tfrac{1}{2})^2}} \end{aligned} \qquad (74)$$

for an emission involving quantum levels n and $n-1$, which becomes

$$\triangle M_n \approx \sqrt{M^2 + \tfrac{q^2}{2} - Qq - \tfrac{1}{2}(n+\tfrac{1}{2})} - \sqrt{M^2 + \tfrac{q^2}{2} - Qq - \tfrac{1}{2}(n-\tfrac{1}{2})} \qquad (75)$$

for large n.

6 Conclusion remarks

We began our paper by summarizing some basic similarities and differences between SBHs and RNBHs in terms of charge and horizon radii. Moreover, we briefly explored the Parikh-Wilczek statement that explains how energy conservation and pair production [2, 21] are fundamentally related to such BHs. For a BH's discrete energy spectrum, the emission or absorption of a particle yields a transition between two distinct levels, where particle emission and absorption are reverse processes [3]-[7]. For this, we touched on the important issue that the nonstrictly thermal character of Hawking's radiation spectrum generates a natural correspondence between Hawking's radiation and BH QNMs, because these structures exemplify features of the BH's energy spectrum [3]-[5], which has been recently generalized to the emerging concept of a BH's effective state [6, 7].

Next, we prepared for our nonextremal RNBH QNM investigation by first reviewing relevant portions of the SBH effective framework [3]-[5] in Section 2. Here, we listed the noneffective and effective quantities for SBH states and transitions, with direct application to the QNM characterization and framework of [3]-[5]. Subsequently, in Section 3, we identified some existing noneffective quantities and introduced new effective quantities for RNBH states and transitions so

we could apply the BH framework of [3]-[5] to implement a RNBH framework. These results are crucial because the effective quantities in [3]-[5] have been achieved for the stable four dimensional RNBH solution in Einstein's general relativity—now effective frameworks exist for the SBH, KBH, and (nonextremal) RNBH solutions.

Ultimately, the RNBH effective quantities permitted us to utilize both the KBH's effective state concept [6, 7] and the BH QNMs [3]-[5] to construct a foundation for the RNBH's effective state in this developing BH effective framework. The RNBH effective state concept is meaningful because, as scientists who wish to demystify the BH paradigm, we need additional features and knowledge to consider in future experiments and observations.

Finally, we stress that the nonstrictly thermal behavior of the Hawking radiation spectrum has been recently used to construct two very intriguing proposals to solve the BH information loss paradox. The first one received the First Award in the 2013 Gravity Research Foundation Essay Competition [26]. The latter won the Community Rating at the 2013 FQXi Essay Contest - It from Bit or Bit from It [28]. We are working to extend this second approach to the RNBH framework [29].

Acknowledgements

S. H. Hendi wishes to thank the Shiraz University Research Council. The work of S. H. Hendi has been supported financially by the Research Institute for Astronomy & Astrophysics of Maragha (RIAAM), Iran.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this article.

References

[1] B. Wang, R. K. Su, P. K. N. Yu and E. C. M. Young, Phys. Rev. D 57, 5284 (1998).

[2] M. K. Parikh and F. Wilczek, Phys. Rev. Lett. 85, 5042 (2000).

[3] C. Corda, JHEP 1108, 101 (2011).

[4] C. Corda, Eur. Phys. J. C 73, 2665 (2013).

[5] C. Corda, Int. Journ. Mod. Phys. D 21, 1242023 (2012, Honorable Mention at Gravity Research Foundation).

[6] C. Corda, S. H. Hendi, R. Katebi, and N. O. Schmidt, JHEP 6, 8 (2013).

[7] C. Corda, Ann. Phys. 337, 49 (2013).

[8] S. Hod, Gen. Rel. Grav. 31, 1639 (1999, Fifth Award at Gravity Research Foundation).

[9] S. Hod, Phys. Rev. Lett. 81 4293 (1998).

[10] M. Maggiore, Phys. Rev. Lett. 100, 141301 (2008).

[11] M. Casals, A. C. Ottewill, Phys. Rev. D 86, 024021 (2012).

[12] T. Padmanabhan, CQG 21, L1 (2004).

[13] A. Lopez-Ortega, CQG 28, 035009 (2011).

[14] S. Hod, CQG 23, L23 (2006).

[15] H. P. Nollert, Phys. Rev. D 47, 5253 (1993).

[16] N. Andersson, CQG 10, L61 (1993).

[17] L. Motl, Adv. Theor. Math. Phys. 6, 1135 (2003).

[18] L. Motl and A. Neitzke, Adv. Theor. Math. Phys. 7, 307 (2003).

[19] S. Hod, Phys. Lett. B 710, 349 (2012).

[20] R. A. Konoplya and A. Zhidenko, Rev. Mod. Phys. 83, 793 (2011).

[21] M. K. Parikh, Gen. Rel. Grav. 36, 2419 (2004, First Award at Gravity Research Foundation).

[22] B. Zhang, Q.-Y. Cai, L. You,, and M. S. Zhan, Phys. Lett. B 675, 98 (2009).

[23] B. Zhang, Q.-Y. Cai, M. S. Zhan, and L. You, Ann. Phys. 326, 350 (2011).

[24] B. Zhang, Q.-Y. Cai, M. S. Zhan, and L. You, EPL 94, 20002 (2011).

[25] B. Zhang, Q.-Y. Cai, M. S. Zhan, and L. You, arXiv:1210.2048.

[26] B. Zhang, Q.-Y. Cai, M. S. Zhan, and L. You, to appear in a Special Issue of Int. Journ. Mod. Phys. D (2013, First Award in the Gravity Research Foundation Essay Competition).

[27] J. Zhang and Z. Zhao, JHEP 10, 55 (2005).

[28] C. Corda, http://fqxi.org/community/forum/topic/1856.

[29] C. Corda, S. H. Hendi, R. Katebi, and N. O. Schmidt, in preparation.

[30] R. Banerjee and B. R. Majhi, Phys. Lett. B 674, 218 (2009).

[31] S. W. Hawking, "The Path Integral Approach to Quantum Gravity", in General Relativity: An Einstein Centenary Survey, eds. S.W.Hawking and W.Israel, (Cambridge University Press, 1979).